

2004-9 Planting List

AREA	SPECIES	PLANTINGS (2004-9)	LOSSES 2005-9	TOTAL (May 2009)
<u>1C</u>				
Plot 12 (10x10m) East Of Dam	Lightwood (<i>Acacia implexa</i>)	2	1	1
	Black Sheoke (<i>Allocasuarina littoralis</i>)	1		1
	Crimson Bottlebrush (<i>Callistemon citrinus</i>)	10	10	0
	Sweet Bursaria (<i>Bursaria spinosa</i>)	2	0	2
	Hedge Wattle (<i>Acacia paradoxa</i>)	2	0	2
Plot 13	NOT PLANTED established for Seed Bank			
Plot 14 (Regeneration Area)	Rough Barked Myrtle (<i>Melaleuca parvistaminea</i>) (1)	30	?	200+*
Scattered Planting	Crimson Bottlebrush (<i>Callistemon citrinus</i>)	7	1	7
	*includes natural regeneration :approximate total only			
<u>2A</u>				
Plot 1 (10mx10m)	Lightwood (<i>Acacia implexa</i>)	3	1	2
	Black Sheoke (<i>Allocasuarina littoralis</i>)	2	0	2
	Sweet Bursaria (<i>Bursaria spinosa</i>)	7	1	6
	Crimson Bottlebrush (<i>Callistemon citrinus</i>)	8	1	7
	Spreading Wattle (<i>Acacia genistifolia</i>)	9	2	7
	Saw Banksia (<i>Banksia serrata</i>)	2	2	0
	Austral Indigo (<i>Indigofera australis</i>)	4	4	0
Plot 2 (10mx10m)	Lightwood (<i>Acacia implexa</i>)	1	0	1
	Black Sheoke (<i>Allocasuarina littoralis</i>)	2	0	2
	Saw Banksia (<i>Banksia serrata</i>)	2	0	2
	Sweet Bursaria (<i>Bursaria spinosa</i>)	6	0	6
	Austral Indigo (<i>Indigofera australis</i>)	4	0	4

AREA	SPECIES	PLANTINGS (2004-9)	LOSSES 2005-9	TOTAL (May 2009)
2A (cont)				
Plot 3 (10mx10m)	Lightwood (<i>Acacia implexa</i>) & Blackwood (<i>Acacia melanoxylon</i>)	4	0	4
	Black Sheoke (<i>Allocasuarina littoralis</i>)	1	0	1
	Saw Banksia (<i>Banksia serrata</i>)	4	1	3
	Sweet Bursaria (<i>Bursaria spinosa</i>)	6	1	5
	Austral Indigo (<i>Indigofera australis</i>)	4	1	3
	Spreading Wattle (<i>Acacia genistifolia</i>)	3	0	3
Plot 4 (10mx12m)	Lightwood (<i>Acacia implexa</i>) & Blackwood (<i>Acacia melanoxylon</i>)	4	1	3
	Black Sheoke (<i>Allocasuarina littoralis</i>)	24	0	24
Plot 5 (12mx14m)	Saw Banksia (<i>Banksia serrata</i>)	22	21	1
	Silver Banksia (<i>Banksia marginata</i>)	11	7	4
	Blackwood (<i>Acacia melanoxylon</i>)	1	0	1
Plot 6 (TFN 10mx10m)	Hop Bitter Pea (<i>Daviesia latifolia</i>)	5	2	3
Plot 7 (TFN 10mx10m)	Spreading Wattle (<i>Acacia genistifolia</i>)	8	7	1
	Heath Wattle (<i>Acacia brownii</i>)	2	0	2
	Snowy Daisy (<i>Olearia lirata</i>)	5	0	5
Plot 8 (TFN 10mx10m)	Spreading Wattle (<i>Acacia genistifolia</i>)	8	4	4
	Hop Bitter Pea (<i>Daviesia latifolia</i>)	2	0	2
	Grey Parrot Pea (<i>Dillwynia cinerascens</i>)	3	2	1
	Snowy Daisy (<i>Olearia lirata</i>)	5	0	5

AREA	SPECIES	PLANTINGS (2004-9)	LOSSES 2005-9	TOTAL (May 2009)
2A (cont)				
Plot 9 (10mx10m)	Heath Wattle (<i>Acacia brownii</i>)	9	6	3
	Snowy Daisy (<i>Olearia lirata</i>)	5	1	4
	Grey Parrot Pea (<i>Dillwynia cinerascens</i>)	8	3	5
	Hop Bitter Pea (<i>Daviesia latifolia</i>)	2	0	2
Plot 10 (10mx12m)	Austral Indigo (<i>Indigofera australis</i>)	15	0	15
	Heath Wattle (<i>Acacia brownii</i>)	4	3	1
	Grey Parrot Pea (<i>Dillwynia cinerascens</i>)	8	7	1
	Hop Bitter Pea (<i>Daviesia latifolia</i>)	2	0	2
	Trigger Plant (<i>Stylidium ameria</i>)	3	?	?
Plot 11 (12mx12m)	NOT PLANTED established for Seed Bank			
Scattered Planting	Saw Banksia (<i>Banksia serrata</i>)	58	35	23
	Black Sheoke (<i>Allocasuarina littoralis</i>)	20	3	17
	Spreading Wattle (<i>Acacia genistifolia</i>)	12	2	10
	Austral Indigo (<i>Indigofera australis</i>)	3	2	1
	Bushy Needlewood (<i>Hakea decurrens</i>)	14	4	10
	Silver Banksia (<i>Banksia marginata</i>)	56	25	31
	Tree Violet (<i>Hymenanthera dentata</i>)	14	7	7
	Drooping Sheoke (<i>Allocasuarina verticillata</i>)	3	0	3
	Lightwood (<i>Acacia implexa</i>)	11	2	9
	Hedge Wattle (<i>Acacia paradoxa</i>)	3	0	3
	Hop Bitter Pea (<i>Daviesia latifolia</i>)	3	1	2
	Purple Diuris (<i>Diuris punctata</i>)	3	?	?
Wedding Bush (<i>Ricinocarpos pinifolius</i>)	2	2	0	

AREA	SPECIES	PLANTINGS (2004-9)	LOSSES 2005-9	TOTAL (May 2009)
<u>2B</u>				
Scattered Planting	Saw Banksia (<i>Banksia serrata</i>)	13	6	7
	Silver Banksia (<i>Banksia marginata</i>)	9	7	2
<u>3A</u>				
Scattered Planting (Regeneration Area)	Lightwood (<i>Acacia implexa</i>) & Blackwood (<i>Acacia melanoxylon</i>)	11	6	5
	Black Sheoke (<i>Allocasuarina littoralis</i>)	16	6	10
	Bushy Needlewood (<i>Hakea decurrens</i>)	6	1	5
	Sweet Bursaria (<i>Bursaria spinosa</i>)	10	3	7
	Hedge Wattle (<i>Acacia paradoxa</i>)	80	22	58
	Crimson Bottlebrush (<i>Callistemon citrinus</i>)	19	6	13
	Spreading Wattle (<i>Acacia genistifolia</i>)	8	6	2
	Tree Violet (<i>Hymenanthera dentata</i>)	8	8	0
TOTALS (2)	(3)	634	245	383

- (1) Plantings and natural regeneration of Rough Barked Myrtle (*Melaleuca parvistaminea*) are not included in final totals.
- (2) Equates to approx 60% survival rate. Removal of Saw Banksia (*Banksia serrata*) and Silver Banksia (*Banksia marginata*) from totals, the two species which suffered excessively high loss rates, increases the survival percentage to 68%.
- (3) Two lots of plantings carried out by TfN in three Exclusion Plots are not included in totals.